arXiv:1110.4297v1 [math.DG] 19 Oct 2011

ON THE CLASSIFICATION OF THE ALMOST CONTACT METRIC MANIFOLDS

VALENTIN A. ALEXIEV AND GEORGI T. GANCHEV

Abstract. The vector space of the tensors $\mathcal{F}$ of type (0,3) having the same symmetries as the covariant derivative of the fundamental form of an almost contact metric manifold is considered. A scheme of decomposition of $\mathcal{F}$ into orthogonal components which are invariant under the action of $U(n)\times 1$ is given. Using this decomposition there are found 12 natural basic classes of almost contact metric manifolds. The classes of cosymplectic, α-Sasakian, α-Kenmotsu, etc. manifolds fit nicely to these considerations. On the other hand, many new interesting classes of almost contact metric manifolds arise.

1. Preliminaries

Let V be a $(2n+1)$-dimensional real vector space with almost contact metric structure (φ,ξ,η,g), where φ is a tensor of type (1,1), ξ is a vector, η is a covector and g is a definite metric so that

$$\varphi^2 x = -x + \eta(x)\,\xi, \quad \varphi(\xi) = 0, \quad \eta\circ\varphi = 0,$$
$$g(\xi,\xi) = 1, \quad g(\varphi x,\varphi y) = g(x,y) - \eta(x)\eta(y)$$

for arbitrary vectors x, y in V. For arbitrary $x \in V$, we denote $hx = -\varphi^2 x$.

We consider the subspace $\mathcal{F}$ of $V^*\otimes V^*\otimes V^*$ defined by the conditions:

$$(1)\ \mathcal{F} = \{F\in\mathcal{F}\,|\,F(x,y,z) = -F(x,z,y) = -F(x,\varphi y,\varphi z) + \eta(y)F(x,\xi,z) + \eta(z)F(x,y,\xi)\}$$

for x,y,z in V.

Let $\{e_i\}, i = 1,...,2n+1$ be an orthonormal basis of V. The metric g induces an inner product in the vector space $\mathcal{F}$:

$$< F', F'' >= \sum_{i,j,k=1}^{2n+1} F'(e_i,e_j,e_k)F''(e_i,e_j,e_k); \quad F', F'' \in \mathcal{F}.$$

We associate with every $F\in\mathcal{F}$ the following covectors:

$$(2)\ f(F)(z) = \sum_i F(e_i,e_i,z), \quad f^*(F)(z) = \sum_i F(e_i,\varphi\, e_i,z), \quad \omega(F)(z) = F(\xi,\xi,z).$$

The standard representation of $U(n)\times 1$ in V induces an associated representation of $U(n)\times 1$ in $\mathcal{F}$.

It is well known the following

Lemma 1. Let L be an involutive isometry of $\mathcal{F}$, which commutes with the action of $U(n)\times 1$ in $\mathcal{F}$. Then

$$\mathcal{F} = L^+(\mathcal{F}) \oplus L^-(\mathcal{F}),$$

2000 *Mathematics Subject Classification.* Primary 53D15, Secondary 53C25.

Key words and phrases. Almost contact metric manifold; covariant derivative of the fundamental form; decomposition of a space of tensors with symmetries.

where $L^+(\mathcal{F})$ and $L^-(\mathcal{F})$ are the eigenspaces of L corresponding to the eigenvalues $+1$ and -1 of L. The decomposition is orthogonal and invariant under the action of $U(n)\times 1$. The components of an element $F\in\mathcal{F}$ in $L^+(\mathcal{F})$ and $L^-(\mathcal{F})$ are

$$F^+=\frac{1}{2}(F+LF),\quad F^-=\frac{1}{2}(L-LF).$$

2. Associated forms with an element of $\mathcal{F}$

With every F in $\mathcal{F}$ we associate the following basic forms:

$$\begin{aligned}
F_1(F)(x,y,z)= &\ \eta(x)F(\xi,y,z),\\
F_2(F)(x,y,z)= &\ \eta(y)F(x,\xi,z)-\eta(z)F(x,\xi,y),\\
F_3(F)(x,y,z)= &\ \eta(x)\eta(y)F(\xi,\xi,z)-\eta(x)\eta(z)F(\xi,\xi,y)\\
= &\ \eta(x)\eta(y)\omega(F)(z)-\eta(x)\eta(z)\omega(F)(y),\\
hF(x,y,z)= &\ F(hx,hy,hz).
\end{aligned}$$

Lemma 2. Let $F\in\mathcal{F}$. Then $F_i(F)(i=1,2,3)$ and hF are elements of $\mathcal{F}$ and

$$F=hF+F_1(F)+F_2(F)-F_3(F).$$

Further we consider the forms

$$\begin{aligned}
&F_4(F)(x,y,z)=\eta(y)F(\varphi x,\xi,\varphi z)-\eta(z)F(\varphi x,\xi,\varphi y),\\
&F_5(F)(x,y,z)=\eta(y)F(z,\xi,x)-\eta(z)F(y,\xi,x),\\
&F_6(F)(x,y,z)=\eta(y)F(\varphi z,\xi,\varphi x)-\eta(z)F(\varphi y,\xi,\varphi x),\\
&F_7(F)(x,y,z)=\frac{1}{2n}f(F)(\xi)\{\eta(z)g(x,y)-\eta(y)g(x,z)\},\\
&F_8(F)(x,y,z)=-\frac{1}{2n}f^*(F)(\xi)\{\eta(z)g(x,\varphi y)-\eta(y)g(x,\varphi z)\}
\end{aligned}$$

associated with an arbitrary $F\in\mathcal{F}$.

Lemma 3. Let $F\in\mathcal{F}$. Then $F_i(F)\,(i=4,5,6,7,8)$ are elements of $\mathcal{F}$.

Lemma 4. Let $F\in\mathcal{F}$. The following relations are valid

$$\begin{aligned}
&F_{11}(F)=F_1(F),F_{12}(F)=F_3(F),F_{13}(F)=F_3(F),F_{14}(F)=0,F_{15}(F)=0,\\
&F_{21}(F)=F_3(F),F_{22}(F)=F_2(F),F_{23}(F)=F_3(F),F_{24}(F)=F_4(F),F_{25}(F)=F_5(F),\\
&F_{31}(F)=F_3(F),F_{32}(F)=F_3(F),F_{33}(F)=F_3(F),F_{34}(F)=0,F_{35}(F)=0,\\
&F_{41}(F)=0,F_{42}(F)=F_4(F),F_{43}(F)=0,F_{44}(F)=F_2(F)-F_3(F),F_{45}(F)=F_6(F),\\
&F_{51}(F)=0,F_{52}(F)=F_5(F),F_{53}(F)=0,F_{54}(F)=F_6(F),F_{55}(F)=F_2(F)-F_3(F),\\
&F_{71}(F)=F_{17}(F)=F_{73}(F)=F_{37}(F)=0,F_{7i}(F)=F_{i7}(F)=F_7(F),\ i=2,4,5,7,\\
&F_{81}(F)=F_{18}(F)=F_{83}(F)=0,F_{58}(F)=F_{85}(F)=-F_{48}(F)=-F_{84}(F)=F_{88}(F)=F_8(F),\\
&h(F_i(F)=F_i(hF)=0,\ i=1,...,8,
\end{aligned}$$

where $F_{ij}(F)=F_i(F_j(F))$.

Lemma 5. Let $F \in \mathcal{F}$. Then we have

$$\begin{aligned}
&f(F_1(F)) = \omega(F),\ f^*(F_1(F)) = 0,\ \omega(F_1(F)) = \omega(F),\\
&f(F_2(F)) = \omega(F) + f(F)(\xi)\eta,\ f^*(F_2(F)) = f^*(F)(\xi)\eta,\ \omega(F_2(F)) = \omega(F),\\
&f(F_3(F)) = \omega(F),\ f^*(F_3(F)) = 0,\ \omega(F_3(F)) = \omega(F),\\
&f(F_4(F)) = f(F)(\xi)\eta,\ f^*(F_4(F)) = f^*(F)(\xi)\eta,\ \omega(F_4(F)) = 0,\\
&f(F_5(F)) = f(F)(\xi)\eta,\ f^*(F_5(F)) = -f^*(F)(\xi)\eta,\ \omega(F_5)(F) = 0,\\
&f(F_6(F)) = f(F)(\xi)\eta,\ f^*(F_6(F)) = -f^*(F)(\xi)\eta,\ \omega(F_6(F)) = 0,\\
&f(F_7(F)) = f(F)(\xi)\eta,\ f^*(F_7(F)) = 0,\ \omega(F_7(F)) = 0,\\
&f(F_8(F)) = 0,\ f^*(F_8(F) = f^*(F)(\xi)\eta,\ \omega(F_8(F)) = 0.
\end{aligned}$$

3. THE SUBSPACES $h\mathcal{F}$, $v\mathcal{F}$ AND $\mathcal{F}_1$ OF $\mathcal{F}$

The first operator L_1. Let $F \in \mathcal{F} :\ L_1(F) = F - 2F_3(F)$.
By straightforward computations, using lemmas 4 and 5, we obtain

Lemma 6. L_1 is an involutive isometry of $\mathcal{F}$ and commutes with the action of $U(n) \times 1$.

This lemma and Lemma 1 imply immediately

Lemma 7. $\mathcal{F}_1 \oplus \mathcal{F}_1^{\perp}$, where

$$\begin{aligned}
\mathcal{F}_1 &= L_1^-(\mathcal{F}) = \{F \in \mathcal{F} \,|\, F = F_3(F)\},\\
\mathcal{F}_1^{\perp} &= L_1^+(\mathcal{F}) = \{F \in \mathcal{F} \,|\, \omega(F) = 0\}.
\end{aligned}$$

The second operator L_2. Let $F \in \mathcal{F}_1^{\perp} :\ L_2(F) = F - 2\{F_1(F) + F_2(F)\}$.
Analogously to Lemma 6 we obtain

Lemma 8. L_2 is an involutive isometry of $\mathcal{F}_1^{\perp}$ and commutes with the action of $U(n) \times 1$. We have

$$\mathcal{F}_1^{\perp} = v\mathcal{F} \oplus h\mathcal{F} \quad \text{(orthogonally)},$$

where

$$\begin{aligned}
v\mathcal{F} &= L_2^-(\mathcal{F}_1^{\perp}) = \{F \in \mathcal{F} \,|\, hF = 0, \omega(F) = 0\},\\
h\mathcal{F} &= L_2^+(\mathcal{F}_1^{\perp}) = \{F \in \mathcal{F} \,|\, F_1(F) = F_2(F) = 0\}.
\end{aligned}$$

Taking into account lemmas 6, 7 and 8, we obtain a partial decomposition:

Proposition 1. $\mathcal{F} = \mathcal{F}_1 \oplus v\mathcal{F} \oplus h\mathcal{F}$. The decomposition is orthogonal and invariant under the action of $U(n) \times 1$. The corresponding components of $F \in \mathcal{F}$ are

$$p_1(F) = F_3(F), \quad vF = F_1(F) + F_2(F) - 2F_3(F), \quad hF.$$

4. THE SUBSPACE $v\mathcal{F}$ OF $\mathcal{F}$

The operator L_3. Let $F \in v\mathcal{F} :\ L_3(F) = F_2(F) - F_1(F)$.

Lemma 9. L_3 is an involutive isometry of $v\mathcal{F}$ and commutes with the action of $U(n) \times 1$. We have $v\mathcal{F} = \mathcal{F}_8 \oplus (v\mathcal{F})'$, where

$$\begin{aligned}
\mathcal{F}_8 &= L_3^-(v\mathcal{F}) = \{F \in \mathcal{F} \,|\, hF = 0, F(x, y, \xi) = 0\},\\
(v\mathcal{F})' = \mathcal{F}_8^{\perp} &= L_3^+(v\mathcal{F}) = \{F \in \mathcal{F} \,|\, hF = 0, F(\xi, y, z) = 0\}.
\end{aligned}$$

The corresponding components of $F \in v\mathcal{F}$ are $F_1(F)$ and $F_2(F)$.

The operator L_4. Let $F \in \mathcal{F}_8^{\perp} = (v\mathcal{F})' :\ L_4(F) = -F_4(F)$.

Lemma 10. L_4 is an involutive isometry of $(v\mathcal{F})' = \mathcal{F}_8^{\perp}$ and commutes with the action of $U(n) \times 1$. We have

$$(v\mathcal{F})' = \mathcal{F}_8^{\perp} = \mathcal{NF}_8^{\perp} \oplus \widetilde{\mathcal{N}}\mathcal{F}_8^{\perp},$$

where

$$\begin{aligned} \mathcal{NF}_8^{\perp} &= L_4^-(\mathcal{F}_8^{\perp}) = \{F \in \mathcal{F} \,|\, F = F_4(F)\}, \\ \widetilde{\mathcal{N}}\mathcal{F}_8^{\perp} &= L_4^+(\mathcal{F}_8^{\perp}) = \{F \in \mathcal{F} \,|\, F = -F_4(F)\}. \end{aligned}$$

The corresponding components of $F \in (v\mathcal{F})' = \mathcal{F}_8^{\perp}$ are

$$\frac{1}{2}\{F_2(F) + F_4(F)\}, \quad \frac{1}{2}\{F_2(F) - F_4(F)\}.$$

The operator L_5. Let $F \in \mathcal{NF}_8^{\perp}$ $(F \in \widetilde{\mathcal{N}}\mathcal{F}_8^{\perp}) : L_5(F) = -F_5(F)$.

Lemma 11. L_5 is an involutive isometry of $\mathcal{NF}_8^{\perp}$ $(\widetilde{\mathcal{N}}\mathcal{F}_8^{\perp})$ and commutes with the action of $U(n) \times 1$. We have

$$\mathcal{NF}_8^{\perp} = \mathcal{QSF} \oplus \mathcal{QKF}, \quad \widetilde{\mathcal{N}}\mathcal{F}_8^{\perp} = \mathcal{F}_6 \oplus \mathcal{F}_7,$$

where

$$\begin{aligned} \mathcal{QSF} &= L_5^-(\mathcal{NF}_8^{\perp}) = \{F \in \mathcal{F} \,|\, F = F_4(F) = F_5(F)\}, \\ \mathcal{QKF} &= L_5^+(\mathcal{NF}_8^{\perp}) = \{F \in \mathcal{F} \,|\, F = F_4(F) = -F_5(F)\}, \\ \mathcal{F}_6 &= L_5^-(\widetilde{\mathcal{N}}\mathcal{F}_8^{\perp}) = \{F \in \mathcal{F} \,|\, F = -F_4(F) = F_5(F)\}, \\ \mathcal{F}_7 &= L_5^+(\widetilde{\mathcal{N}}\mathcal{F}_8^{\perp}) = \{F \in \mathcal{F} \,|\, F = -F_4(F) = -F_5(F)\}. \end{aligned}$$

The corresponding components of $F \in \mathcal{NF}_8^{\perp}$ $(F \in \widetilde{\mathcal{N}}\mathcal{F}_8^{\perp})$ are

$$\begin{aligned} &\frac{1}{4}\{F_2(F) + F_4(F) + F_5(F) + F_6(F)\}, \quad \frac{1}{4}\{F_2(F) + F_4(F) - F_5(F) - F_6(F)\}, \\ &\left(\frac{1}{4}\{F_2(F) - F_4(F) + F_5(F) - F_6(F)\}, \quad \frac{1}{4}\{F_2(F) - F_4(F) - F_5(F) + F_6(F)\}\right). \end{aligned}$$

The operator L_6. Let $F \in \mathcal{QSF} :\ L_6(F) = F - 2F_7(F)$.

Lemma 12. L_6 is an involutive isometry of $\mathcal{QSF}$ and commutes with the action of $U(n) \times 1$. We have

$$\mathcal{QSF} = \mathcal{F}_2 \oplus \mathcal{F}_4 \quad \text{(orthogonally)},$$

where

$$\begin{aligned} \mathcal{F}_2 &= L_6^-(\mathcal{QSF}) = \{F \in \mathcal{F} \,|\, F = F_7(F)\}, \\ \mathcal{F}_4 &= L_6^+(\mathcal{QSF}) = \{F \in \mathcal{F} \,|\, F = F_4(F) = F_5(F),\ f(F)(\xi) = 0\}. \end{aligned}$$

The corresponding components of $F \in \mathcal{QSF}$ in $\mathcal{F}_2$ and $\mathcal{F}_4$ are

$$F_7(F), \quad \frac{1}{4}\{F_2(F) + F_4(F) + F_5(F) + F_(6) - 4F_7(F)\}.$$

The operator L_7. Let $F \in \mathcal{QKF} :\ L_7(F) = F - 2F_8(F)$.

Lemma 13. L_7 is an involutive isometry of $\mathcal{QKF}$ and commutes with the action of $U(n) \times 1$. We have

$$\mathcal{QKF} = \mathcal{F}_3 \oplus F_5 \quad \text{(orthogonally)},$$

where

$$\begin{aligned}
&\mathcal{F}_3 = L_7^-(\mathcal{QKF}) = \{F \in \mathcal{F} \,|\, F = F_8(F)\},\\
&\mathcal{F}_5 = L_7^+(\mathcal{QKF}) = \{F \in \mathcal{F} \,|\, F = F_4(F) = -F_5(F),\ f^*(F)(\xi) = 0\}.
\end{aligned}$$

The corresponding components of $F \in \mathcal{QKF}$ in $\mathcal{F}_3$ and $\mathcal{F}_5$ are

$$F_8(F), \quad \frac{1}{4}\{F_2(F) + F_4(F) - F_5(F) - F_6(F) - 4F_8(F)\}.$$

Using lemmas 9 - 13, we get

Proposition 2. $vF = \mathcal{F}_2 \oplus ... \oplus \mathcal{F}_8$. The decomposition is orthogonal and invariant under the action of $U(n) \times 1$. The corresponding components of $F \in \mathcal{F}$ in $\mathcal{F}_i\,(i = 2, ..., 8)$ are

$$\begin{aligned}
&p_2(F) = F_7(F),\\
&p_3(F) = F_8(F),\\
&p_4(F) = \frac{1}{4}\{F_2(F) + F_4(F) + F_5(F) + F_6(F) - 4F_7(F) - F_3(F)\},\\
&p_5(F) = \frac{1}{4}\{F_2(F) + F_4(F) - F_5(F) - F_6(F) - 4F_8(F) - F_3(F)\},\\
&p_6(F) = \frac{1}{4}\{F_2(F) - F_4(F) + F_5(F) - F_6(F) - F_3(F)\},\\
&p_7(F) = \frac{1}{4}\{F_2(F) - F_4(F) - F_5(F) + F_6(F) - F_3(F)\},\\
&p_8(F) = F_1(F) - F_3(F).
\end{aligned}$$

5. The subspace $h\mathcal{F}$

Now, let $hV = \{x \in V \,|\, x = hx\}$. Denoting the restrictions of g and φ on hV with the same letters, we obtain the Hermitian vector space $\{hV, g, \varphi\}$ of dimension $2n$. We identify the elements of $h\mathcal{F}$ with their restrictions on hV. Then we can consider the vector space $h\mathcal{F}$ as the vector space of the tensors hF of type (0,3) over hV having the properties

$$hF(x, y, z) = -hF(x, z, y) = -hF(x, \varphi y, \varphi z)$$

for all $x, y, z \in hV$. The action of $U(n) \times 1$ on $h\mathcal{F}$ coincides with the action of $U(n)$ on $h\mathcal{F}$. In [1] the vector space $h\mathcal{F}$ has been decomposed orthogonally into irreducible components invariant under the action of $U(n)$.

Let $F \in h\mathcal{F}$. It is not difficult to verify that the forms

$$\begin{aligned}
F_9(F) = \frac{1}{2(n-1)} \quad &\{g(hx, hy)f(F)(z) - g(hx, hz)f(F)(y)\\
&-g(x, \varphi y)f(F)(\varphi z) + g(x, \varphi z)f(F)(\varphi y)\},\\
F_{10}(F) = \frac{1}{2}\,&\{F(x, y, z) + f(\varphi x, \varphi y, z)\},\\
F_{11}(F) = \frac{1}{6} \quad &\{F(x, y, z) + F(y, z, x) + F(z, x, y)\\
&-F(\varphi x, \varphi y, z) - F(\varphi y, \varphi z, x) - F(\varphi z, \varphi x, y)\},\\
F_{12}(F) = \frac{1}{2}\,&\{F(x, y, z) - F(\varphi x, \varphi y, z)\},
\end{aligned}$$

are also elements of $h\mathcal{F}$.

Using the decomposition in [1], we have

Proposition 3. $h\mathcal{F} = F_9 \oplus F_{10} \oplus F_{11} \oplus F_{11} \oplus F_{12}$, where

$$\mathcal{F}_9 = \{F \in \mathcal{F} \,|\, F = hF = F_9(F)\},$$
$$\mathcal{F}_{10} = \{F \in \mathcal{F} \,|\, F = hF = F_{10}(F) - F_9(F)\},$$
$$\mathcal{F}_{11} = \{F \in \mathcal{F} \,|\, F = hF = F_{11}(F)\},$$
$$\mathcal{F}_{12} = \{F \in \mathcal{F} \,|\, F = hF = F_{12}(F) - F_{11}(F)\}.$$

The decomposition is orthogonal and invariant under the action of $U(n) \times 1$. The corresponding components of $F \in \mathcal{F}$ are

$$F_9(F), \quad F_{10}(F) - F_9(F), \quad F_{11}(F), \quad F_{12}(F) - F_{11}(F).$$

6. APPLICATIONS TO ALMOST CONTACT METRIC MANIFOLDS

Let M be an almost contact metric manifold with structure (φ, ξ, η, g), where φ is a tensor field of type (1,1), ξ is a tensor field, η is a 1-form, and g is a Riemannian metric on M such that

$$\varphi^2 x = -x + \eta(x)\,\xi, \quad g(\xi,\xi) = 1, \quad \eta \circ \varphi = 0,$$
$$\varphi\,\xi = 0, \quad g(\varphi x, \varphi y) = g(x,y) - \eta(x)\eta(y)$$

for arbitrary vector fields x, y on M. For all vector fields x, y, z on M we denote

$$F(x,y,z) = g((\nabla_x \varphi)y, z). \tag{3}$$

Let T_pM be the tangent space to M at $p \in M$ and $V = T_pM$. The restriction F_p of F given by (3) on V has the properties (1). We shall call M is of class $\mathcal{W}_i$ $(i = 1, ..., 12)$ if F_p is in the subspace $\mathcal{F}_i$ $(i = 1, ..., 12)$ for every $p \in M$. Using the propositions 1, 2 and 3 we obtain 12 basic classes of almost contact metric manifolds . Further we give the defining conditions for these classes. Let F be given by (3) and f, f^*, ω be $f(F), f^*(F), \omega(F)$ respectively defined by (2).

The class $\mathcal{W}_1$*:*

$$F(x,y,z) = \eta(x)\eta(y)\omega(z) - \eta(x)\eta(z)\omega(y).$$

The class $\mathcal{W}_2$*:*

$$F(x,y,z) = \frac{f(\xi)}{2n}\{\eta(z)g(x,y) - \eta(y)g(x,z)\}.$$

This is the class of α-Sasakian manifolds.

The class $\mathcal{W}_3$*:*

$$F(x,y,z) = -\frac{f^*(\xi)}{2n}\{\eta(z)g(x,\varphi y) - \eta(y)g(x,\varphi z)\}.$$

This is the class of α-Kenmotsu manifolds.

The class $\mathcal{W}_4$*:*

$$\begin{aligned} F(x,y,z) &= \eta(y)F(\varphi x, \xi, \varphi z) - \eta(z)F(\varphi x, \xi, \varphi y) \\ &= \eta(y)F(z,\xi,x) - \eta(z)F(y,\xi,x), \qquad f(\xi) = 0. \end{aligned}$$

The class $\mathcal{W}_5$*:*

$$\begin{aligned} F(x,y,z) &= \eta(y)F(\varphi x,\xi,\varphi z) - \eta(z)F(\varphi x,\xi,\varphi y) \\ &= -\eta(y)F(z,\xi,x) + \eta(z)F(y,\xi,x), \qquad f^*(\xi) = 0. \end{aligned}$$

The class $\mathcal{W}_6$*:*

$$\begin{aligned} F(x,y,z) &= -\eta(y)F(\varphi x,\xi,\varphi z) + \eta(z)F(\varphi x,\xi,\varphi y) \\ &= \eta(y)F(z,\xi,x) - \eta(z)F(y,\xi,x). \end{aligned}$$

The class $\mathcal{W}_7$*:*

$$\begin{aligned} F(x,y,z) &= -\eta(y)F(\varphi x,\xi,\varphi z) + \eta(z)F(\varphi x,\xi,\varphi y) \\ &= -\eta(y)F(z,\xi,x) + \eta(z)F(y,\xi,x). \end{aligned}$$

The class $\mathcal{W}_8$*:*

$$F(hx,hy,hz) = F(x,y,\xi) = 0.$$

The class $\mathcal{W}_9$*:*

$$\begin{gathered} F(\xi,y,z) = F(x,y,\xi) = 0, \\ F(x,y,z) = \frac{1}{2(n-1)} \begin{aligned}[t] &\{g(\varphi x,\varphi y)f(z) - g(\varphi x,\varphi z)f(y) \\ &-g(x,\varphi y)f(\varphi z) + g(x,\varphi z)f(\varphi y)\}. \end{aligned} \end{gathered}$$

The class $\mathcal{W}_{10}$*:*

$$\begin{gathered} F(\xi,y,z) = F(x,y,\xi) = 0, \\ F(\varphi x,\varphi y,z) - F(x,y,z) = 0, \quad f = 0. \end{gathered}$$

The class $\mathcal{W}_{11}$*:*

$$\begin{gathered} F(\xi,y,z) = F(x,y,\xi) = 0, \\ F(x,x,z) = 0. \end{gathered}$$

The class $\mathcal{W}_{12}$*:*

$$\begin{gathered} F(\xi,y,z) = F(x,y,\xi) = 0, \\ F(x,y,z) + F(y,z,x) + F(z,x,y) = 0. \end{gathered}$$

The class of cosymplectic manifolds is characterized by $F = 0$. This class is contained in all $\mathcal{W}_i$ $(i = 1,...,12)$. An almost contact metric manifold M belongs to two classes $\mathcal{W}_i$, $\mathcal{W}_j$ $(i \neq j)$ iff M is cosymplectic.

References

[1] Gray A. and L. Hervella. *The Sixteen Classes of Almost Hermitian Manifolds and Their Linear Invariants.* Ann. Mat. Pura Appl. **123** (1980), 35-58.

Bulgarian Academy of Sciences, Institute of Mathematics and Informatics, Acad. G. Bonchev Str. bl. 8, 1113 Sofia, Bulgaria
E-mail address: `ganchev@math.bas.bg`